後発医薬品の使用に伴い生じたインシデント・アクシデント事例の 実態とその解析

堀里子¹)、寺門千佳子²)、石河利恒²)、大和田邦明²)、字野幹子¹)、三木晶子¹)、大谷壽一¹)、 天野武城²)、澤田康文^{1,3)*}

¹⁾ 東京大学大学院薬学系研究科 医薬品情報学講座:東京都文京区本郷 7-3-1

2) 東邦薬品株式会社:東京都世田谷区代沢 5-2-1

¹⁾ 東京大学大学院情報学環・学際情報学府:東京都文京区本郷 7-3-1

An Investigation on Cases of Accidents and Incidents Related to Use of Generic Drugs

Satoko HORI¹), Chikako TERAKADO²), Toshitsune ISHIKO²), Kuniaki OHWADA²), Mikiko UNO¹), Akiko MIKI¹), Hisakazu OHTANI¹), Takeshiro AMANO²) and Yasufumi SAWADA^{1,3}*

¹⁾ Laboratory of Drug Informatics, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

²⁾ Toho Pharmacentical CO., LTD. 5-2-1 Daizawa, Setagaya-ku, Tokyo 155-8655, Japan

³⁾ Graduate School of Interdisciplinary Information Studies, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

(Recevied August 20, 2007) Accepted November 2, 2007)

Abstract

We collected cases of accidents and incidents related to the use of generic drugs from pharmacies and medical facilities, and analyzed their causes. Eighty nine troubles were collected, and most of them occurred when prescription was switched from the branded formulation to their generic counterparts. The most frequent trouble was the confusion of medicines (49 cases), most of which are confusion of branded drug and generic one within same constituent and same content. The other cases resulted from the differences in the indications between branded and generic drugs (11 cases), the differences in the feel of use or usability between branded and generic drugs (9 cases), and the duplicate prescription of drugs with same constituents or similar effects (3 cases).

It would be important for pharmacists to report the accidents and incidents related the use of to genetic drugs, and share safety measures against them. This would lead to promoting proper use of genetic drugs by preventing medical errors related to the use of genetic drugs.

Key words: generic drugs, accidents and incidents